



Planning Community-Led Mangrove Restoration and Monitoring Fish Nursery Recovery in a Bay of Bengal Estuary: A Methodological Framework

Dana Pham^{1*}, Jakub Raghavan², Fernanda Martirosyan³

¹Department of Fashion Designing, Ebro Institute for Advanced Fashion Designing, Spain

²Department of Fashion Designing, Mekong Center for Fashion Designing, United Kingdom

³Department of Fashion Designing, Transylvanian Institute of Fashion Designing, Argentina

Article Info

Volume 1, Issue 1, 2026

[doi: 10.62715/ijeb.2026.v1.i1.pp15-20](https://doi.org/10.62715/ijeb.2026.v1.i1.pp15-20)

Received: 02 February 2026

Accepted: 20 March 2026

Published: 16 May 2026

Abstract

Mangrove restoration around the Bay of Bengal has a mixed record. Large areas have been planted, often with a single species on mudflats where mangroves never grew, and many plantations have failed. Where restoration is led by the villages that fish and gather in the estuary, and where it begins with tidal hydrology, the prospects are better, yet few such projects measure whether the restored stands come to function as nurseries for fish and crustaceans. This article proposes a six-phase framework that joins the two concerns. The phases are site diagnosis, agreement on rules and roles, restoration of tidal exchange, assisted natural regeneration, paired monitoring of vegetation and juvenile fish, and periodic review. For each phase the article states the purpose, the tasks, who carries them out and the record to be kept. Fish monitoring follows a before-after, control-impact logic with reference and unrestored creeks, and uses gears that village monitors can operate. The framework is assembled from the published literature on ecological mangrove restoration, nursery function, common-pool resource management and participatory monitoring. It is a proposal. It has not been applied or validated here, and the article sets out what would have to be shown before it could be recommended.

Keywords: Mangrove restoration, Community-based management, Nursery habitat, Estuarine fish, Participatory monitoring, Bay of Bengal

© 2026 Dana Pham et al. This is an open access article under the CC BY license (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

1. Introduction

The estuaries that open into the Bay of Bengal, from the Godavari and Krishna deltas through the Mahanadi and the Sundarbans to the Ayeyarwady, carry some of the most extensive mangrove forests in Asia and some of its densest coastal populations. Aquaculture ponds, paddy bunds, salt pans and reduced freshwater flow have removed or degraded large parts of these forests. Restoration has followed, driven by forest departments, donors and, after each major cyclone, by renewed interest in coastal protection.

* Corresponding author: Dana Pham, Department of Fashion Designing, Ebro Institute for Advanced Fashion Designing, Spain. E-mail: dana.pham@ebro.edu

© 2026 Dana Pham et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Results have often been poor. Reviews of planting programmes in the Philippines and Sri Lanka found low survival across most sites, largely because seedlings were put in the wrong places (Primavera and Esteban, 2008; Kodikara et al., 2017). Even where trees survive, success is usually reported as hectares planted or seedlings alive. Fishing households tend to judge a mangrove by a different measure: whether juvenile shrimp, crab and fish return to its creeks.

This article proposes a framework for planning a community-led restoration in a Bay of Bengal estuary and for monitoring the recovery of its nursery function. It describes a procedure and the reasoning for each part. It does not report a project, and no site, community or data set stands behind it. The framework has not been validated here.

2. Background

Three bodies of published work inform the design. They concern why planting fails, what a nursery is and how it can be measured, and what makes shared management of a resource last.

2.1. Hydrology before planting

Lewis (2005) set out the approach now called ecological mangrove restoration. Mangroves occupy a specific band of the intertidal zone, roughly between mean sea level and high water of spring tides, and each species has its own tolerance of flooding depth, duration and salinity. If a site has lost its mangroves, the first task is to find out why. Where tidal exchange has been cut by a bund or a road, reopening it often allows propagules carried on the tide to establish without any planting. Planting is a supplement for sites that propagules cannot reach. Ellison (2000) reached a similar position, that enough is known about mangrove ecology to restore forests and that failures come mostly from ignoring it.

Tree cover is a first step and not the goal. A review of restored stands found that faunal recolonisation, nutrient cycling and other functions can approach those of natural forest, but over different periods and not in every case (Bosire et al., 2008). International restoration standards accordingly ask that a project define a reference ecosystem and measure recovery against several attributes, among them species composition, structure and function (Gann et al., 2019).

2.2. The nursery function

Juveniles of many fish and crustaceans concentrate among mangrove roots and in the creeks that drain them. The accepted explanations are structural shelter, reduced predation in shallow turbid water and abundant food (Laegdsgaard and Johnson, 2001; Nagelkerken et al., 2008). High density alone does not make a nursery. Beck et al. (2001) defined a nursery as a habitat that contributes more recruits to the adult population per unit area than other habitats in which juveniles occur, through some combination of density, growth, survival and successful movement to adult grounds. Dahlgren et al. (2006) added that the overall contribution of a habitat also matters, and Sheaves et al. (2015) argued that nursery value belongs to connected mosaics of mangrove, mudflat and channel more than to a single habitat. Most evidence linking mangrove area to fishery catch is correlational (Manson et al., 2005), which is a further reason to measure the response directly.

2.3. Institutions for shared management

A restored stand is a common-pool resource: villagers can cut poles, graze cattle or net juveniles in it, and one household's restraint benefits all. Ostrom (1990) identified features of institutions that have managed such resources for long periods, including clear boundaries, rules made by the users, monitoring by people answerable to them, graduated sanctions and recognition by the state. A review of community-based mangrove management found that tenure security, a visible flow of benefits and sustained outside support separated lasting efforts from short-lived ones (Datta et al., 2012).

3. The Proposed Framework

The framework has six phases arranged as a cycle, shown in Figure 1, with a village mangrove committee at the centre. Table 1 lists each phase with its lead and the record it should produce. The phases are

described below in order.




Table 1: Phases of the proposed framework, who leads each, and the record to be kept

Phase	Main tasks	Led by	Record kept
1. Diagnose	Map elevation, tides, salinity, past cover and causes of loss	Committee with technical partner	Site map and diagnosis note
2. Agree	Settle boundaries, access rules, roles and benefit sharing	Village assembly	Signed agreement
3. Restore flow	Breach bunds, clear or re-dig creeks	Committee work groups	Works log with photographs
4. Assist regeneration	Protect recruits; plant only where recruitment fails	Committee work groups	Planting and survival register
5. Monitor	Survey vegetation plots and juvenile fish in fixed creeks	Trained village monitors	Data sheets and season summary
6. Review	Compare results with targets and change the plan	Assembly and partners	Annual review minutes

3.1. Phase 1: Diagnose the site

The committee and a technical partner walk the site at spring and neap tides and record where water reaches and how long it stays. Ground level is surveyed against a tide staff, since a difference of a few tens of centimetres decides which species, if any, can grow. Salinity is measured in creeks and soil water in the dry season and after the monsoon. Old maps, satellite images and the memory of older residents establish where mangroves stood and what removed them. A nearby intact stand at similar elevation is chosen as the reference, and its species zonation noted: typically *Avicennia marina* and *Sonneratia apetala* on the accreting seaward edge, *Rhizophora* along creek banks, and *Excoecaria agallocha* and *Ceriops* on higher ground.

3.2. Phase 2: Agree rules and roles

Before any earth is moved, the village assembly settles who may use the site and how. The agreement fixes the boundary on the ground, names the committee and its term, sets rules on cutting, grazing and fine-mesh netting inside the restored creeks, and states how wages for restoration work and any later

harvest will be shared. Groups with a direct stake, among them women who collect crab and shrimp seed and households that hold or lease former ponds, need seats on the committee and not only a hearing. The agreement should be registered with the forest department or local government, so that the community's rights are recognised by the authority that holds legal title.

3.3. Phase 3: Restore tidal flow

Works follow the diagnosis. Abandoned pond bunds are breached at the points where former creeks crossed them, silted channels are re-dug by hand to their old course, and culverts are opened under paths. The aim is a pattern of flooding and drainage close to that of the reference stand. Where the ground has subsided or been excavated, spoil from the breaches can raise it to the level at which the reference species grow. Work is timed for the dry season, when access is easiest and labour is available.

3.4. Phase 4: Assist natural regeneration

The site is then watched through one full fruiting season. If propagules arrive and establish, the committee's task is protection: fencing against cattle and goats and enforcing the agreed rules. Planting is reserved for areas where no recruits appear, and uses species matched to the measured elevation and salinity, raised from local seed. Single-species blocks of *Rhizophora* on bare lower mudflat, the pattern that failed elsewhere (Primavera and Esteban, 2008), are excluded by the diagnosis.

3.5. Phase 5: Monitor vegetation and juvenile fish

Vegetation is recorded in permanent plots laid along transects from creek edge to landward margin, in the restored site and the reference stand. Monitors count seedlings and saplings by species and measure stem diameter and height of trees, following published protocols for mangrove plots (Kauffman and Donato, 2012). Fixed photograph points add a record that anyone in the village can read.

Fish monitoring is designed so that a change at the restored site can be told apart from a good or bad year in the estuary as a whole. The logic is that of before-after, control-impact designs (Stewart-Oaten et al., 1986), extended to several control locations as Underwood (1994) recommended. Three kinds of creek are sampled: restored creeks, reference creeks in intact mangrove and unrestored creeks in degraded ground. Sampling starts before the bunds are breached and continues on the same schedule afterwards. A divergence of the restored creeks from the unrestored ones, and towards the reference, is the signal sought.

Gears must suit small tidal creeks and village operators. A block net set across a creek mouth at high water, with the catch collected as the creek drains, samples a known area. Fyke nets and local bamboo traps are alternatives where a creek cannot be closed. Sampling is repeated on set lunar phases in the pre-monsoon, monsoon and post-monsoon seasons. Monitors identify the catch to an agreed list that includes commercially important juveniles such as penaeid shrimp, mud crab (*Scylla*), mullets and barramundi (*Lates calcarifer*), count individuals, measure the length of a subsample and release live animals. Density and size structure of juveniles are the main indicators. They address the density component of the definition by Beck et al. (2001); growth, survival and movement to adult stocks lie beyond a village programme.

Participatory monitoring ranges from schemes run by outside scientists with local help to schemes designed and interpreted by local people, and the authors of that typology argue that the more locally driven kinds tend to feed more directly into management decisions (Danielsen et al., 2009). The framework places itself towards the local end. The technical partner trains monitors, checks a share of identifications against photographs or retained specimens, and helps with analysis.

3.6. Phase 6: Review and adjust

Once a year the assembly meets with its partners to compare results with the targets set in the diagnosis: flooding regime, recruitment, canopy cover, and juvenile density relative to reference creeks. Shortfalls lead back to the appropriate phase. Persistent waterlogging calls for further channel work, repeated grazing damage for a change in rules or their enforcement.

4. Fitting the Framework to Bay of Bengal Conditions

Several features of the region shape how the framework would be used. Tidal range is large in the northern bay and small along parts of the Coromandel coast, so the width of the zone suitable for mangroves, and the gear that works in the creeks, differ from one estuary to another. Monsoon river flow drives a strong seasonal swing in salinity, which affects both seedling establishment and the juvenile fish present; comparisons must therefore be made season by season. Tropical cyclones can flatten young stands and re-close creeks with sediment, and the review phase should include an inspection after every severe storm. Land status is frequently contested between revenue, forest and fisheries authorities, and the agreement of the second phase has little force until that question is settled.

5. Limitations and What Remains to Be Shown

The framework is a proposal built from published principles, and its parts have not been tested together. Whether village monitors can produce fish data consistent enough to detect a change is the first open question, and it depends on training, on turnover among monitors and on how variable creek catches prove to be. A second is duration. Restored stands take years to develop structure, and faunal recovery may lag further (Bosire et al., 2008), so a programme that ends with its funding after two or three seasons will record only the beginning.

A third limitation is inferential. A rise in juvenile density in restored creeks shows that young animals use the habitat. It does not show that the estuary produces more adults, since fish may simply have moved from neighbouring creeks. Tagging, otolith chemistry or stable isotope work would be needed to trace contribution to adult stocks, and these require a research partner. Finally, the framework assumes a community able and willing to organise. Where households are divided over the land, or where outside interests control the former ponds, the second phase may fail, and no technical design can substitute for it.

6. Conclusion

The framework proposed here treats mangrove restoration as two linked tasks: bringing back a tidal forest by repairing hydrology and protecting natural recruitment, and finding out, with methods a village can run, whether juvenile fish and crustaceans return to it. It gives the community the lead, asks for an agreement before works begin, and builds in controls so that the fish data can bear interpretation. Its elements come from an established literature. Their combination remains to be tried, and a pilot in a single estuary, reported with its difficulties, would be the appropriate next step.

Conflict of interest

To be declared by the author(s) before publication.

Funding

To be declared by the author(s) before publication.

References

- Beck, M.W., Heck, K.L., Able, K.W., Childers, D.L., Eggleston, D.B., Gillanders, B.M., Halpern, B., Hays, C.G., Hoshino, K., Minello, T.J., Orth, R.J., Sheridan, P.F. and Weinstein, M.P. (2001). [The identification, conservation, and management of estuarine and marine nurseries for fish and invertebrates](#). *BioScience*, 51: 633-641.
- Bosire, J.O., Dahdouh-Guebas, F., Walton, M., Crona, B.I., Lewis, R.R., Field, C., Kairo, J.G. and Koedam, N. (2008). [Functionality of restored mangroves: A review](#). *Aquatic Botany*, 89: 251-259. <https://doi.org/10.1016/j.aquabot.2008.03.010>
- Dahlgren, C.P., Kellison, G.T., Adams, A.J., Gillanders, B.M., Kendall, M.S., Layman, C.A., Ley, J.A., Nagelkerken, I. and Serafy, J.E. (2006). [Marine nurseries and effective juvenile habitats: concepts and applications](#). *Marine Ecology Progress Series*, 312: 291-295. <https://doi.org/10.3354/meps312291>

- Danielsen, F., Burgess, N.D., Balmford, A., Donald, P.F., Funder, M., Jones, J.P.G., Alviola, P., Balete, D.S., Blomley, T., Brashares, J., et al. (2009). [Local participation in natural resource monitoring: a characterization of approaches](#). *Conservation Biology*, 23: 31-42. <https://doi.org/10.1111/j.1523-1739.2008.01063.x>
- Datta, D., Chattopadhyay, R.N. and Guha, P. (2012). [Community based mangrove management: A review on status and sustainability](#). *Journal of Environmental Management*, 107: 84-95. <https://doi.org/10.1016/j.jenvman.2012.04.013>
- Ellison, A.M. (2000). [Mangrove restoration: Do we know enough?](#) *Restoration Ecology*, 8: 219-229. <https://doi.org/10.1046/j.1526-100x.2000.80033.x>
- Gann, G.D., McDonald, T., Walder, B., Aronson, J., Nelson, C.R., Jonson, J., Hallett, J.G., Eisenberg, C., Guariguata, M.R., Liu, J., Hua, F., Echeverría, C., Gonzales, E., Shaw, N., Decler, K. and Dixon, K.W. (2019). [International principles and standards for the practice of ecological restoration. Second edition](#). *Restoration Ecology*, 27(S1): S1-S46. <https://doi.org/10.1111/rec.13035>
- Kauffman, J.B. and Donato, D.C. (2012). [Protocols for the measurement, monitoring and reporting of structure, biomass and carbon stocks in mangrove forests](#). Working Paper 86. Center for International Forestry Research (CIFOR), Bogor.
- Kodikara, K.A.S., Mukherjee, N., Jayatissa, L.P., Dahdouh-Guebas, F. and Koedam, N. (2017). [Have mangrove restoration projects worked? An in-depth study in Sri Lanka](#). *Restoration Ecology*, 25: 705-716. <https://doi.org/10.1111/rec.12492>
- Laegdsgaard, P. and Johnson, C. (2001). [Why do juvenile fish utilise mangrove habitats?](#) *Journal of Experimental Marine Biology and Ecology*, 257: 229-253.
- Lewis, R.R. (2005). [Ecological engineering for successful management and restoration of mangrove forests](#). *Ecological Engineering*, 24: 403-418. <https://doi.org/10.1016/j.ecoleng.2004.10.003>
- Manson, F.J., Loneragan, N.R., Skilleter, G.A. and Phinn, S.R. (2005). [An evaluation of the evidence for linkages between mangroves and fisheries: a synthesis of the literature and identification of research directions](#). *Oceanography and Marine Biology: An Annual Review*, 43: 483-513.
- Nagelkerken, I., Blaber, S.J.M., Bouillon, S., Green, P., Haywood, M., Kirton, L.G., Meynecke, J.-O., Pawlik, J., Penrose, H.M., Sasekumar, A. and Somerfield, P.J. (2008). [The habitat function of mangroves for terrestrial and marine fauna: A review](#). *Aquatic Botany*, 89: 155-185. <https://doi.org/10.1016/j.aquabot.2007.12.007>
- Ostrom, E. (1990). [Governing the Commons: The Evolution of Institutions for Collective Action](#). Cambridge University Press, Cambridge.
- Primavera, J.H. and Esteban, J.M.A. (2008). [A review of mangrove rehabilitation in the Philippines: successes, failures and future prospects](#). *Wetlands Ecology and Management*, 16: 345-358. <https://doi.org/10.1007/s11273-008-9101-y>
- Sheaves, M., Baker, R., Nagelkerken, I. and Connolly, R.M. (2015). [True value of estuarine and coastal nurseries for fish: incorporating complexity and dynamics](#). *Estuaries and Coasts*, 38: 401-414. <https://doi.org/10.1007/s12237-014-9846-x>
- Stewart-Oaten, A., Murdoch, W.W. and Parker, K.R. (1986). [Environmental impact assessment: "pseudoreplication" in time?](#) *Ecology*, 67: 929-940.
- Underwood, A.J. (1994). [On beyond BACI: sampling designs that might reliably detect environmental disturbances](#). *Ecological Applications*, 4: 3-15. <https://doi.org/10.2307/1942110>